



## 2.1 Bisection Method (Binary Search)

Play High/Low.





### Algorithm

Input  $a, b, \text{TOL}, \text{max \# of iterations } N_0$

Output approx solution  $p$  or message of failure

Step 1 for  $(i=1 \text{ to } N_0)$  do step 2-4.

Step 2 set  $p = \frac{a+b}{2}$

Step 3 if  $f(p) = 0$  or  $\underbrace{(b-a)/2 < \text{TOL}}$  then  
can do other too.

output( $p$ );  
Stop

Step 4 If  $f(a)f(p) > 0$  then set  $a = p$   
else set  $b = p$

Step 5 Output ('Method failed after  $N_0$  iteration!') Stop.

Try on  $f(x) = x^2 - 2x + 1 = 0$ .

**Thm 2.1**

Let  $f \in C[a, b]$  and suppose  $f(a) \cdot f(b) < 0$ .

The Bisection method (Alg. 2.1) generates a sequence  $\{P_n\}$  approximating  $p$  with the property


$$|P_n - p| \leq \frac{b-a}{2^n}, \quad n \geq 1$$

Proof:  $\forall n \geq 1$ , we have

$$b_n - a_n = \frac{1}{2^{n-1}} (b-a)$$

(We half the interval each step.)

Note that  $p \in (a_n, b_n)$ .



Since  $P_n = \frac{1}{2}(a_n + b_n)$ ,  $\forall n \geq 1$ , it follows that

$$\frac{1}{2}(b-a) \quad b-a$$

$$|p_n - p| \leq \frac{1}{2} (b_n - a_n) = \frac{1}{2} \frac{2^{n-1}}{2^{n-1}} = \frac{1}{2^n}$$

Thus, as  $n \rightarrow \infty$ ,  $p_n \rightarrow p$  at the rate of  $O\left(\frac{1}{2^n}\right)$ .

---

Ex: How many iterations of Bisection would be required to get the approx accurate to within  $10^{-8}$   
Suppose  $a=0, b=1$ . Then

$$|p_n - p| \leq \frac{b-a}{2^n} < 10^{-8}$$

$$2^{-n} < 10^{-8}$$

$$-n \log 2 < \log 10^{-8}$$

$$n > \frac{8}{\log 2} = 26.57 \Rightarrow 27 \text{ iterations}$$